



Chile (Region VIII)

2004 Fact Sheet

The **2004 Chile (Region VIII) GSHS** measured alcohol and other drug use; sexual behaviors that contribute to HIV infection, other STI, and unintended pregnancy; unintentional injuries and violence; hygiene; dietary behaviors and overweight; physical activity; tobacco use; mental health; and protective factors.

The 2004 Chile (Region VIII) GSHS was a school-based survey of students in grades 7° primary, 8° primary, and 1° secondary. A two-stage cluster sample design was used to produce data representative of all students in grades 7° primary, 8° primary, and 1° secondary in Chile (Region VIII). At the first stage, schools were selected with probability proportional to enrollment size. At the second stage, classes were randomly selected and all students in selected classes were eligible to participate. The school response rate was 100%, the student response rate was 87%, and the overall response rate was 87%. A total of 1,971 students participated in the Chile (Region VIII) GSHS. Students self-reported their responses to each question on a computer scannable answer sheet. For comparison purposes, only students aged 13-15 years are included in the analyses for this fact sheet.

Results for students aged 13-15 years

Total

Boys

Girls

Alcohol and Other Drug Use

Percentage of students who had at least one drink containing alcohol on one or more days during the past 30 days	25.4 ± 3.2	24.8 ± 4.9	25.9 ± 3.8
Percentage of students who drank so much alcohol that they were really drunk one or more times during their life	21.9 ± 4.5	23.1 ± 5.0	20.3 ± 4.7
Percentage of students who had a hang-over, felt sick, got into trouble with their family or friends, missed school, or got into fights, as a result of drinking alcohol one or more times during their life	10.3 ± 3.0	10.8 ± 3.5	9.6 ± 3.2
Percentage of students who used drugs one or more times during their life	7.2 ± 2.0	7.8 ± 2.5	6.3 ± 2.3

Sexual Behaviors That Contribute to HIV Infection, Other STI, and Unintended Pregnancy

Percentage of students who have ever had sexual intercourse	13.0 ± 3.0	18.9 ± 3.9	7.5 ± 2.9
Percentage of students who have had sexual intercourse with two or more people during their life	5.7 ± 1.8	9.0 ± 3.1	2.6 ± 0.9
Among students who had sexual intercourse during the past 12 months, the percentage who used a condom the last time they had sexual intercourse	51.5 ± 10.9	*	*

Unintentional Injuries and Violence

Percentage of students who were in a physical fight one or more times during the past 12 months	34.0 ± 3.4	49.0 ± 4.6	19.2 ± 3.0
Percentage of students who were seriously injured one or more times during the past 12 months	37.0 ± 4.5	43.9 ± 5.8	30.4 ± 4.5
Percentage of students who were bullied on one or more days during the past 30 days	49.8 ± 4.6	53.6 ± 4.8	46.3 ± 6.1



Chile (Region VIII)

2004 Fact Sheet

Results for students aged 13-15 years	Total	Boys	Girls
Hygiene			
Percentage of students who never or rarely washed their hands before eating during the past 30 days	2.8 ± 0.9	2.6 ± 1.3	2.9 ± 1.4
Percentage of students who never or rarely washed their hands after using the toilet or latrine during the past 30 days	1.1 ± 0.7	1.3 ± 1.0	0.8 ± 0.7
Dietary Behaviors and Overweight			
Percentage of students who went hungry most of the time or always during the past 30 days because there was not enough food in their home	1.1 ± 0.5	0.5 ± 0.5	1.6 ± 0.8
Percentage of students who are overweight**	1.9 ± 0.9	2.3 ± 1.1	1.5 ± 1.3
Percentage of students who are at risk for becoming overweight***	15.5 ± 2.1	15.0 ± 3.1	16.0 ± 3.1
Physical Activity			
Percentage of students who were physically active for a total of at least 60 minutes per day on all 7 days during the past 7 days	8.9 ± 2.0	12.4 ± 2.8	5.4 ± 2.3
Percentage of students who spent three or more hours per day during a typical or usual day sitting and watching television, playing computer games, talking with friends, or doing other sitting activities	40.1 ± 4.4	31.9 ± 4.1	48.0 ± 5.9
Tobacco Use			
Percentage of students who smoked cigarettes on one or more days during the past 30 days	23.0 ± 4.3	24.2 ± 5.4	21.9 ± 4.5
Percentage of students who used any other form of tobacco on one or more days during the past 30 days	4.1 ± 1.4	4.4 ± 2.4	3.7 ± 1.5
Percentage of students who reported people smoking in their presence on one or more days during the past 7 days	76.5 ± 2.1	78.0 ± 4.1	75.1 ± 3.3
Mental Health			
Percentage of students who felt lonely most of the time or always during the past 12 months	14.1 ± 1.8	7.7 ± 2.1	20.1 ± 3.1
Percentage of students who seriously considered attempting suicide during the past 12 months	19.5 ± 2.5	11.6 ± 4.0	26.9 ± 3.2
Percentage of students who have no close friends	6.8 ± 1.3	7.3 ± 1.8	6.3 ± 1.5



Chile (Region VIII) 2004 Fact Sheet

Results for students aged 13-15 years	Total	Boys	Girls
Protective Factors			
Percentage of students who missed classes or school without permission on one or more days during the past 30 days	17.9 ± 3.1	20.8 ± 4.5	15.1 ± 3.7
Percentage of students who reported that most of the students in their school were never or rarely kind and helpful during the past 30 days	30.3 ± 2.6	33.1 ± 3.4	27.4 ± 3.8
Percentage of students whose parents or guardians never or rarely really knew what they were doing with their free time during the past 30 days	27.7 ± 3.4	28.3 ± 4.4	27.0 ± 4.9

*Indicates data were not available.

**Students who were at or above the 95th percentile for body mass index by age and sex based on reference data from Cole, Bellizzi, Flegal, and Dietz, British Medical Journal, May, 2000.

***Students who were at or above the 85th percentile but below the 95th percentile for body mass index by age and sex based on reference data from Cole, Bellizzi, Flegal, and Dietz, British Medical Journal, May, 2000.

For additional information, please contact:

Claudia Gonzalez
Ministerio de Salud de Chile, Santiago, Chile, cgonzale@minsal.cl